

F-7241

METHOD FOR DETERMINING SIGNIFICANT BONE DENSITY LOSSES

The invention relates to a method for determining significant bone density losses.

In the interaction of bone destruction processes and bone formation processes, there may, on the average, be loss of bone substance and structure. In order to determine this loss and identify significant deviations from age-specific and sex-specific reference values, bone density measurements (osteodensitometry) are generally carried out. Using the methods of photon adsorption or computer tomography, the bone mineral density is measured at the radius, at the thigh bone (femur) or at the lumbar vertebral column (Vertebra lumbalis). Under the assumption of an exponential development of the bone density decrease, the loss can be estimated from bone density values measured at least at three different times. Intra-individual measurement fluctuations of up to 5 percent may be obtained. In order to keep these factors, which interfere with the determination of the results, as small as possible, sufficiently large intervals between the times of measuring the bone density should be selected, so that changes in the bone density can be detected adequately. At the earliest, therefore, information is available after one year.

Osteodensitometric methods of identifying bone density losses are expensive, cost intensive and expose the patient to radiation. Moreover, osteodensitometry is not transferable between different instruments and measurement methods are not standardized. At the present time, there are no recognized alternative methods with equivalent informative capability.

In the literature (P. Garnero and P.D. Dalmás, "Biochemical Markers of Bone Turnover" 1988, Endocrinology and Metabolism Clinics of North America, vol.

27, No. 2, pp. 303 -322), laboratory parameters, which are particularly associated with bone density losses (bone markers), are investigated extensively with respect to their suitability for describing bone density loss quantitatively. Such parameters from the cell portion (osteoblasts), from the organic matrix of the bone (collagen, noncollageous proteins) or from the inorganic constituents of the bone admittedly can give an indication of bone loss. Until now, however, the information, provided by them, is not equivalent to the information provided by osteodensitometry. With that, the advantage of simplicity of this method of determining parameters in the serum or the urine cannot be utilized.

It is an object of the invention to develop a method for determining significant bone density losses, which is less cost intensive, does not expose the patient to radiation and shortens the period for interpreting the interaction of bone destruction processes and bone formation processes.

Pursuant to the invention, this objective is accomplished owing to the fact that measurement values of real or mathematically simulated bone density loss processes, which are present in electronic storage media and, as a function of time, reflect laboratory parameters of practically or theoretically known clinical signs and symptoms, are used as reference values for the process, that bone marker values of serum or urine samples, associated with bone density losses and measured by common laboratory techniques, are determined by sample preparation steps such as

- treating with antibodies
- incubation steps
- separation procedures
- using analytical techniques

and recorded over an input mask in an electronic data memory, are used for determining significant bone density losses,

- a) at the time of the analysis, all N available patient-related data being copied from the data memory over an interrogation function and made available for

10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532

- c) the normalized measured value being converted into a scalar quantity $D(t_n)$ for the graduated description of the course of the bone density, the equation

$$D(t_n) = \sqrt{\sum_{k=1}^K W_k \cdot (M^*(t_n; k))^2}$$

being used as functions for the graduated description of the course of the relationship

- $$D^*(t) = \frac{(t_n - t) \cdot D(n-1) + (t - t_{n-1}) \cdot D(n)}{t_n - t_{n-1}}, t \in [t_{n-1}, t_n]$$

3

- e) from the interpolated evaluations of the progress, similarity dimensions being calculated, the function

$$A_j(t) = \sum_{m=1}^M \frac{t_m}{t_M} \cdot V_m \cdot (R_j(t_m) - D^*(t_m))^2$$

being used to calculate a similarity dimension between the data, which is to be investigated, and all the reference values, available in the database and, at the same time, similarity dimensions to the reference values and to the time in months being found;

- f) from the similarity dimensions for all reference values, those reference values being determined, which have a high similarity in the mathematical sense, such as the

greatest similarity: $A^* = \min_{j=1, \dots, J} \{A_j\}$

positive alternative (+) $A^+ = \min_{j=1, \dots, J, A_j \neq A^*, R_j(t(N)) > D(t(N))} \{A_j\}$

negative alternative (-) $A^- = \min_{j=1, \dots, J, A_j \neq A^*, R_j(t(N)) < D(t(N))} \{A_j\}$

with subsequent output of the type description as text component for describing the situation;

- g) the prediction being derived from these three reference values, if $B_1 = A^*$, $B_2 = A^+$, $B_3 = A^-$,

the following expression

$$R(t) = \frac{1}{\sum_{i=1}^3 B_i} \cdot \sum_{j=1}^3 \left(\left(\sum_{i=1}^3 B_i - B_j \right) \cdot R_j(t) \right)$$

being used for the predicted value at time t

- h) the degrees of freedom for the specification of the model, given as functional parameters in the functional relation of $D(t_n)$ and $A_j(t)$ being occupied by standard specifications and fitted by statistical analysis of the reference values to practical experience for optimizing the quantitative prediction of the bone density loss;
- i) the time being calculated, at which, according to this prediction strategy, the percentage deviation is greater than a specified threshold value, this time being the starting point for planning the scheduling of the next investigation.

Advantageously, the degrees of freedom, given as function parameters in the functional relationship of $D(t_n)$ and $A_j(t)$, are filled in by the mathematical method of least squares so that specified sequences are taken into consideration in the best way for reference values.

The reference values used can be values from an assumed analytical mathematical course (exponential function), from empirically based values from imaginary, assumed processes and from concrete, measured values from patients with known situations.

In the following, the invention is explained in greater detail by means of an example.

Osteocalcin, parathyroid hormone and alkaline phosphatase are used as bone markers. Data for the bone markers are obtained from serum or urine samples using well-known laboratory techniques (HPLC, RIA, ELISA). For this purpose, sample preparation steps such as

- treatment with antibodies
- incubation steps
- separation methods
- use in analytical techniques

are necessary in order to obtain, after the measurement, a quantitative value in the parameter-specific unit as a monitor or printer output or as an electronically available numerical value. These laboratory values obtained are stored in an electronic database over an input mask. It is a prerequisite of the method that reference values are known. Reference values can be values calculated theoretically from an analytically mathematically assumed course (exponential function) or empirically based values from imaginary, assumed processes or concrete values from patients with known situations. These reference values are available for specified times and can be used in the analysis only within the scope of the time horizon determined therewith. In the sample, the exponential function $R(t) = a(1 - e^{-bt})$ is used to describe bone density losses. The parameters a and b then describe the speed and the degree of loss (t in month). Some reference examples are listed in the following Table:

Months After First Measurement	Type 0 a = 0	Type I a = 1 b = 0.01	Type II a = 0.5 b = 0.05	Type III a = 0.5 b = 0.1	Type IV a = 2 b = 0.01	Type V a = 1 b = 0.05	Type VI a = 2 b = 0.05
0	0.00	0.00	0.00	0.00	0.00	0.00	0.00
6	0.00	0.06	0.13	0.23	0.12	0.26	0.52
12	0.00	0.11	0.23	0.35	0.23	0.45	0.90
18	0.00	0.16	0.30	0.42	0.33	0.59	1.19
24	0.00	0.21	0.35	0.45	0.43	0.70	1.40
30	0.00	0.26	0.39	0.48	0.52	0.78	1.55
36	0.00	0.30	0.42	0.49	0.60	0.83	1.67
42	0.00	0.34	0.44	0.49	0.69	0.88	1.76
48	0.00	0.38	0.45	0.50	0.76	0.91	1.82
54	0.00	0.42	0.47	0.50	0.83	0.93	1.87
60	0.00	0.45	0.48	0.50	0.90	0.95	1.90

This is followed by the steps below

- a) the values measured and those obtained on three prior occasions are given in the Table below:

No.	Date	Osteocalcine in $\mu\text{g/L}$	PTH in ng/L	AP in U/L
1	5-30-96	9.8	24.8	90
2	1-29-97	10.9	34.6	86
3	2-16-98	12.6	32.0	104
4	3-2-99	12.4	34.0	107

measured values $M(t_n; k)$ for $n = 1, \dots, 4$ and $k = 1, \dots, 3$

- b) the values obtained by measurement are normalized with respect to the first line in the Table according to the following equation

$$M^*(t_n; k) = \frac{M(t_n; k) - M(t_1; k)}{M(t_1; k)}$$

and the time interval between the measurements is converted into months;

No.	Month	$M^*(t; 1)$	$M^*(t; 2)$	$M^*(t; 3)$
1	0.0	0.00	0.00	0.00
2	8.0	0.11	0.40	-0.04
3	20.6	0.29	0.29	0.16
4	33.1	0.27	0.37	0.19

Normalized values measured $M(t_n; k)$ for $n = 1, \dots, 4$ and $k = 1, \dots, 3$

- c) the normalized measured value is converted into a scalar quantity for the graduated description of the bone density loss, the equation

$$D(t_n) = \sqrt{\sum_{k=1}^K W_k \cdot (M^*(t_n; k))^2}$$

being used ($K = 3; n = 1, \dots, 3$) as a function of the graduated description of the progress. Under the standard setup, it is, of course assumed that $w = 1$ for all weighting factors;

Evaluations of Progress at Time t

- d) From the evaluations of the progress obtained, evaluations of the progress for those time intervals of

$$D^*(t) = \frac{(t_n - t) \cdot D(n-1) + (t - t_{n-1}) \cdot D(n)}{t_n - t_{n-1}}, t \in [t_{n-1}, t_n]$$

are calculated by interpolation, for which reference values are available;

Interpolated Evaluations of Progress at Imaginary Measurement Times
at 6-Month Intervals.

- e) from the interpolated evaluations of the progress, similarity dimensions are calculated, the function

$$A_j(t) = \sum_{m=1}^M \frac{t_m}{t_M} \cdot V_m \cdot (R_j(t_m - D^*(t_m)))^2, j = 1, \dots, 6; M = 6,$$

being used for calculating a similarity dimension between the data, which is to be investigated, and all reference values available in the database, the following similarity dimensions being found. Under a standard setup, $V = 1$ for all weighting factors.

Month		Type 0	Type I	Type II	Type III	Type IV	Type V	Type VI
0								
6	0.32	0.10	0.07	0.04	0.01	0.04	0.00	0.04
12	0.43	0.24	0.14	0.06	0.01	0.06	0.00	0.24
18	0.44	0.36	0.17	0.06	0.01	0.05	0.02	0.72
24	0.47	0.48	0.19	0.06	0.01	0.04	0.07	1.41
30	0.49	0.63	0.21	0.06	0.00	0.03	0.14	2.25

Similarity Dimensions for all Reference Values and all Times up to 30 Months

The instructions for the calculations explicitly do not represent a function for estimating the parameters a and b of the exponential function, since general reference values do not have to satisfy these instructions;

- f) from the similarity dimensions for all reference values, those reference values are determined, which have a high similarity in the mathematical sense, such as the

greatest similarity: $A^* = \min_{j=1, \dots, J} \{A_j\} = 0,00$

positive alternative (+) $A^+ = \min_{j=1, \dots, J, A_j \neq A^*, R_j(N) > D(N)} \{A_j\} = 0,03$

negative alternative (-) $A^- = \min_{j=1, \dots, J, A_j \neq A^*, R_j(N) < D(N)} \{A_j\} = 0,06$

In accordance with these selection instructions, Type III is recognized as the reference value and Type IV and Type II are selected as positive and negative, alternative reference value respectively;

- g) The prediction is derived from these three reference value changes. If, in order to simplify the notation,

$B_1 = A^*$, $B_2 = A^+$, $B_3 = A^-$, the following is used as the predicted value at time t

$$R(t) = \frac{1}{\sum_{i=1}^3 B_i} \cdot \sum_{j=1}^3 \left(\left(\sum_{i=1}^3 B_i - B_j \right) \cdot R_j(t) \right)$$

- h) The degrees of freedom for the specification of the model, given as functional parameters in the functional relation of $D(t_n)$ and $A_j(t)$, are filled in by reference values, in order to achieve a quantitative prediction of the bond density loss;

Month	Prediction for Type III	Prediction for Alternatives (+)	Prediction for Alternatives (-)	Predicted Value $R(t)$	Deviation from Prediction in %
36	0.49	0.60	0.42	0.52	6.1
42	0.49	0.69	0.44	0.55	12.2
48	0.50	0.76	0.45	0.58	16.0
54	0.50	0.83	0.47	0.61	22.0
60	0.50	0.90	0.48	0.63	26.0

Predicted Value

The certainty of this prediction is characterized by the percentage deviation of the average predicted value $R(t)$, using only the suitable reference value, ascertained with the similarity dimension factors A_1 , in the example, prediction for

Type III, which has the greatest similarity A^*

- i) The time is calculated, at which, according to this prediction strategy, the percentage deviation is greater than a specified threshold value, this time being the starting point for planning the scheduling for the subsequent investigation. For this purpose, this value is put out over a monitor or printer and transmitted by remote data transmission to the treating physician. In the case of the present example, the threshold value is assumed with a 10% deviation. Forty months after the first measurement, a deviation greater than 10% is then found for the first time. The time for the next measurement should therefore not be later than 7 months after the values for the bone markers are determined.

Month	Prediction for Type III	Prediction for Alternatives (+)	Prediction for Alternatives (-)	Predicted Value R(t)	Deviation from Prediction in %
40	0.49	0.66	0.43	0.54	10.0

Deviation of the predicted value greater than 10%.